



COLLEGE of AMERICAN
PATHOLOGISTS

ARCHIVES

of Pathology & Laboratory Medicine

EARLY ONLINE RELEASE

This article was posted on the *Archives* Web site as an Early Online Release. Note: Due to the extremely time sensitive nature of the content of this article, it has not been copyedited or formatted per journal style. Changes or corrections may be made to this article when it appears in a future print issue of the *Archives*. Early Online Release articles are citable by using the Digital Object Identifier (DOI), a unique number given to every article.

The DOI for this manuscript is doi: [10.5858/arpa.2021-0435-SA](https://doi.org/10.5858/arpa.2021-0435-SA)

The final published version of this manuscript will replace the Early Online Release version at the above DOI once it is available.

**Autopsy Histopathologic Cardiac Findings in Two Adolescents Following the Second
COVID-19 Vaccine Dose**

James R. Gill, MD; Randy Tashjian, MD; Emily Duncanson, MD

Connecticut Office of the Chief Medical Examiner, Farmington, Connecticut, (Gill); Department of Pathology, Yale School of Medicine, New Haven, Connecticut (Gill); Wayne County Medical Examiners' Office, Detroit, Michigan (Tashjian); Department of Pathology, University of Michigan, Ann Arbor, Michigan (Tashjian); Jesse E. Edwards Registry of Cardiovascular Disease, St. Paul, MN (Duncanson)

Corresponding author:

James Gill, MD

OCME

11 Shuttle Rd

Farmington, CT 06032

jgill@ocme.org

Supplemental digital content can be found at the end of article.

The authors have no relevant financial interest in the products or companies described in this article.

Running title: COVID-19 Heart Vaccine

ABSTRACT

Context.– Myocarditis in adolescents has been diagnosed clinically following the administration of the second dose of an mRNA vaccine for coronavirus disease 2019 (COVID-19).

Objective.– To examine the autopsy microscopic cardiac findings in adolescent deaths that occurred shortly following administration of the second Pfizer-BioNTech COVID-19 dose to determine if the "myocarditis" described in these instances has the typical histopathology of myocarditis.

Design.– Clinical and autopsy investigation of two teenage boys who died shortly following administration of the second Pfizer-BioNTech COVID-19 dose.

Results.– The microscopic examination revealed features resembling a catecholamine-induced injury, not typical myocarditis pathology.

Conclusions.– The myocardial injury seen in these post-vaccine hearts is different from typical myocarditis and has an appearance most closely resembling a catecholamine-mediated stress (toxic) cardiomyopathy. Understanding that these instances are different from typical myocarditis and that cytokine storm has a known feedback loop with catecholamines may help guide screening and therapy.

Myocarditis in adolescents (particularly teenage boys) has been reported following the second dose of the Pfizer-BioNTech COVID-19 vaccine.¹⁻⁷ Since cardiac biopsies are rarely performed in these instances with clinically stable patients, the myocardial pathology has not been clearly elucidated.⁸ Myocarditis is rarely diagnosed at autopsy in deaths due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection.^{9,10} The incidence of myocarditis, although low, has been shown to increase after the receipt of the BNT162b2 vaccine, particularly after the second dose among young male recipients.¹¹ In addition, the first week after the second vaccine dose was found to be the main risk window.¹¹ The clinical presentation of myocarditis after vaccination was usually mild.¹¹

We report the autopsy results including microscopic myocardial findings of two teenage boys who died within the first week after receiving the second Pfizer-BioNTech COVID-19 dose. The microscopic findings are not the alterations seen with typical myocarditis. This suggest a role for cytokine storm which may occur with an excessive inflammatory response, as there also is a feedback loop between catecholamines and cytokines.¹²

MATERIALS AND METHODS

The Connecticut Office of the Chief Medical Examiner (OCME) and the Michigan Institute of Forensic Science and Medicine investigate all unexpected and unnatural deaths in their respective jurisdictions: Connecticut and the Michigan counties of Alcona, Gladwin, Huron, Lapeer, Ogemaw, and Saginaw.

Standard medicolegal autopsies were performed including gross, microscopic, and toxicological testing. SARS-COV-2 nasal swab testing was performed by reverse transcriptase-polymerase chain reaction (RT-PCR) assay. Tissues were sent to the National Center for

Emerging and Zoonotic Infectious Disease Branch of the Centers for Disease Control and Prevention (CDC) for molecular studies.

Cardiac molecular testing with sequence analysis and deletion/duplication testing of the 100 genes listed in Invitae's arrhythmia and cardiomyopathy comprehensive panel was performed.

RESULTS

The results of autopsies for two teenage boys who were found dead in bed 3 and 4 days after receiving the second dose of the Pfizer-BioNTech COVID-19 vaccine are presented (Table 1). Both boys were pronounced dead at home without attempted resuscitation.

Boy A complained of a headache and gastric upset but felt better by post-vaccine day 3. There was no history of prior medical problems (he took prescribed amphetamine/dextroamphetamine during the school year for attention deficit hyperactivity disorder but was not currently receiving it) or prior SARS-COV-2 infection. Boy B had no complaints, prior health issues, or prior SARS-COV-2 infection. Neither boy complained of fever, chest pain, palpitations, or dyspnea. The autopsies were unremarkable except for obesity in one boy and the cardiac findings (Figures 1-7 and Supplemental Figures 1-4 [Supplemental digital content can be found at the end of article]). Unique cardiac findings in Boy A included myocardial fibrosis and in Boy B cardiac hypertrophy. There were no rashes or lymphadenopathy.

Expanded forensic toxicological testing was negative for medications and drugs of abuse. SARS-COV-2 was not detected by postmortem swab (RT-PCR assay) in either boy. Cardiac

sections were submitted from the right and left ventricles (12 sections in Boy A and 29 sections in Boy B). The cardiac conduction systems were not examined.

DISCUSSION

Myocarditis is an inflammatory disease of the myocardium, which may occur in isolation or as part of multiorgan/systemic immune-mediated disorders or reactions to exogenous/endogenous substances.¹³ The etiologies are varied and include infectious and non-infectious causes. Non-infectious causes include immune/autoimmune (autoantigens, association with immune-mediated diseases, alloantigens, and allergens), drugs/toxic substances (e.g., hypersensitivity or direct toxic effects), and other causes (e.g., radiation, and insect stings, snake bites).¹³ Lymphocytic myocarditis is the commonest histological subtype, characterized by an inflammatory myocardial infiltrate typically comprising mononuclear cells. In the acute/active phases, it is usually accompanied by myocyte damage/necrosis.¹³ Although criteria are evolving, the Dallas Criteria requires "inflammatory infiltrates of the myocardium with necrosis and/or degeneration of *adjacent* myocytes, not typical of ischemic damage associated with coronary artery disease."¹⁴⁻¹⁶

Toxic myocarditis is an etiological classification involving direct myocardial injury by various drugs or substances.^{13,17,18} Although variable, the histologic features consist of two main patterns: an early stage with foci of solely necrotic/damaged myocytes and the later phase of "myocarditis." Toxic myocarditis usually indicates inflammatory stages of catecholamine-induced myocardial injury. Catecholamine toxicity on the heart was first described in patients with pheochromocytoma.¹⁹⁻²¹ These lesions have been described in patients with subarachnoid hemorrhages and, more recently, in donor hearts rejected for transplantation in deaths declared

dead by neurological criteria, secondary to catecholamine-release during the ‘sympathetic storm’ following "brain-death" or administered as pharmacologic support (see supplemental images).^{22,23} The wide spectrum of these lesions has been studied in detail in routine pathology examination of donor hearts unsuitable for transplantation.²²

Both teenage boys had similar clinical presentations with no obvious cardiac symptoms. Their histopathology does not demonstrate a typical myocarditis. In those instances, one sees lymphocytic (or giant cell) infiltrates with adjacent myocyte necrosis; changes such as hypereosinophilic myocytes and contraction bands are absent. In these two post-vaccination instances, there are areas of contraction bands and hypereosinophilic myocytes distinct from the inflammation. This injury pattern is instead similar to what is seen in the myocardium of patients who are clinically diagnosed with Takotsubo, toxic, or “stress” cardiomyopathy, which is a temporary myocardial injury that can develop in patients with extreme physical, chemical, or sometimes emotional stressors.²⁴⁻³¹

Stress cardiomyopathy is a catecholamine-mediated ischemic process seen in high catecholamine states in the absence of coronary artery disease or spasm.^{17,31} It has also been called “neurogenic myocardial injury” and “broken heart syndrome.”^{18,24-36} Surges in catecholamines may have several triggers (fight/flight response, adrenal pathology, etc.). Proposed mechanisms for catecholamine-mediated stunning in stress cardiomyopathy include epicardial spasm, microvascular dysfunction, hyperdynamic contractility with midventricular or outflow tract obstruction, and direct effects of catecholamines on cardiomyocytes.³³

Catecholamine-mediated myocardial stunning may be due to direct myocyte injury as elevated catecholamines decrease the viability of myocytes through cyclic adenosine monophosphate (AMP)–mediated calcium overload. Catecholamines also are a potential source

of oxygen-derived free radicals which can interfere with sodium and calcium transporters, possibly resulting in myocyte dysfunction through increased transsarcolemmal calcium influx and cellular calcium overload.³⁷

Histologically, catecholamine effects have been associated with contraction band necrosis, characterized by hypercontracted sarcomeres, dense eosinophilic transverse bands, and an interstitial mononuclear inflammatory response that is distinct from the polymorphonuclear inflammation seen with infarction. In addition, the mononuclear cells are not causing the myocyte necrosis; there is a distinct, separate distribution.³⁷

We suspect that the acute cardiac changes seen in these two boys are the result of epinephrine-mediated effects on cardiomyocytes. These occurrences generally have a favorable prognosis, however, some patients may die from the underlying (non-cardiac) cause of the myocardial findings (e.g., such as with subarachnoid hemorrhage). Histologically, diffuse hyper eosinophilic myocytes, contraction bands, and coagulative myocytolysis are seen, with a patchy and random pattern and a neutrophilic/mononuclear cell infiltrate. With longer survival, global myocardial ischemia may develop.³⁷

This post-vaccine reaction may represent an overly exuberant immune response and the myocardial injury is mediated by similar immune mechanisms as described with SARS-COV-2 and multisystem inflammatory syndrome (MIS-C) cytokine storms.³⁸ MIS-C is a rare systemic illness presenting with persistent fever and extreme inflammation following exposure to SARS-CoV-2. Affected children have persistent fever and may have acute abdominal pain with diarrhea or vomiting, muscle pain/malaise, and hypotension. Other reported symptoms include rashes, enlarged lymph nodes, and swelling.

A hypersensitivity reaction is in the differential diagnosis, however, infrequency/lack of eosinophils would be unusual. The common denominator of a hypersensitivity reaction is the eosinophilic infiltrate, which may be the major inflammatory component or be part of a complex picture of mixed inflammation with lymphocytes, macrophages, plasma cells, poorly formed microgranulomas, and giant cells.³⁹ An autopsy study of 69 cases of hypersensitivity myocarditis examined the spectrum of histologic findings including the distribution of infiltrates and the extent and composition of the infiltrates.⁴⁰ They reported that hypersensitivity myocarditis was "defined by the presence of eosinophils, a mixed lymphohistiocytic infiltrate along natural planes of separation, and an absence of fibrosis or granulation tissue in areas of infiltrate."⁴⁰

Despite a molecular investigation, the etiology of the fibrosis in case A is unclear. It is conceivable that this process first started with the first vaccination dose and the initial myocardial effects resolved and healed over time. The second dose may have restarted the process. One might expect some scarring/repair after a few weeks, although the scarring in Case A appears more organized than the three-week interval between the vaccine doses. Also, it is only in one of the cases. It remains possible that the fibrosis represents arrhythmogenic cardiomyopathy. Unfortunately, cardiac molecular testing was equivocal.

Regardless of the etiology of the fibrosis, the extent of scarring by itself is potentially arrhythmogenic and may be a contributing factor with the acute post-vaccine myocardial injury. Similarly, the cardiac hypertrophy in Case B may have made the heart more susceptible to an arrhythmia. The key point is that since these boys had died suddenly and unexpectedly in their sleep without resuscitation, if the arrhythmia had been due to the myocardial scar (Boy A) or cardiomegaly (Boy B), then the fulminant, global myocardial injury would not be an expected

finding. These two clinical histories support the etiology of the acute myocardial injury as a primary factor not a secondary agonal or post-resuscitative artefact.

Two adults (ages 42 and 45 years) with "myocarditis" diagnosed histologically (one at autopsy and one by biopsy) following SARS-COV-2 mRNA vaccinations were recently reported.⁴¹ One occurred 10 days after receiving the first Pfizer-BioNTech COVID-19 vaccine dose and the other occurred 14 days after receiving the second mRNA-1273 (Moderna) dose. Histologically, both were described as "fulminant" myocarditis with "multifocal cardiomyocyte damage associated with mixed inflammatory infiltration." In addition to areas of myocyte necrosis associated with the inflammatory infiltrate, the photomicrographs demonstrate ischemic changes distinct from the inflammation similar to our findings.

Cytokine storm has been described with an excessive and uncontrolled inflammatory response, and there is a feedback loop between catecholamines and cytokines.¹² Clinical complications may include cardiac compromise, respiratory distress, and hypercoagulation.⁴² The myocardial injury seen in these post-vaccine hearts has a similar histologic appearance as catecholamine-mediated stress cardiomyopathy and severe SARS-COV-2 infection, including "myocarditis" which is associated with cytokine release syndrome.³⁸ Recognition that these instances are different from typical myocarditis and that cytokine storm has a known feedback loop with catecholamines may help guide screening, diagnosis, and therapy.

REFERENCES

1. Wise J. Covid-19: Should we be worried about reports of myocarditis and pericarditis after mRNA vaccines? *BMJ*. 2021;373:n1635.
2. Shay DK, Shimabukuro TT, DeStefano F. Myocarditis Occurring After Immunization With mRNA-Based COVID-19 Vaccines. *JAMA Cardiol*. 2021;6(10):1115-1117.
3. Gargano JW, Wallace M, Hadler SC, et al. Use of mRNA COVID-19 Vaccine After Reports of Myocarditis Among Vaccine Recipients: Update from the Advisory Committee on Immunization Practices - United States, June 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(27):977-982.
4. Rosner CM, Genovese L, Tehrani BN, et al. Myocarditis Temporally Associated With COVID-19 Vaccination. *Circulation*. 2021;144(6):502-505.
5. Navar AM, McNally E, Yancy CW, O'Gara PT, Bonow RO. Temporal Associations Between Immunization With the COVID-19 mRNA Vaccines and Myocarditis: The Vaccine Safety Surveillance System Is Working. *JAMA Cardiol*. 2021;6(10):1117-1118.
6. Muthukumar A, Narasimhan M, Li QZ, et al. In-Depth Evaluation of a Case of Presumed Myocarditis After the Second Dose of COVID-19 mRNA Vaccine. *Circulation*. 2021;144(6):487-498.
7. Montgomery J, Ryan M, Engler R, et al. Myocarditis Following Immunization With mRNA COVID-19 Vaccines in Members of the US Military. *JAMA Cardiol*. 2021;6(10):1202-1206.
8. Marshall M, Ferguson ID, Lewis P, et al. Symptomatic Acute Myocarditis in 7 Adolescents After Pfizer-BioNTech COVID-19 Vaccination. *Pediatrics*. 2021;148(3).

9. Halushka MK, Vander Heide RS. Myocarditis is rare in COVID-19 autopsies: cardiovascular findings across 277 postmortem examinations. *Cardiovasc Pathol.* 2021;50:107300.
10. Fox SE, Falgout L, Vander Heide RS. COVID-19 myocarditis: quantitative analysis of the inflammatory infiltrate and a proposed mechanism. *Cardiovasc Pathol.* 2021;54:107361.
11. Mevorach D, Anis E, Cedar N, et al. Myocarditis after BNT162b2 mRNA Vaccine against Covid-19 in Israel. *N Engl J Med.* 2021;385(23):2140-2149.
12. Staedtke V, Bai RY, Kim K, et al. Disruption of a self-amplifying catecholamine loop reduces cytokine release syndrome. *Nature.* 2018;564(7735):273-277.
13. Leone O, Pieroni M, Rapezzi C, Olivotto I. The spectrum of myocarditis: from pathology to the clinics. *Virchows Arch.* 2019;475(3):279-301.
14. Aretz HT. Myocarditis: the Dallas criteria. *Hum Pathol.* 1987;18(6):619-624.
15. Baughman KL. Diagnosis of myocarditis: death of Dallas criteria. *Circulation.* 2006;113(4):593-595.
16. Basso C, Calabrese F, Angelini A, Carturan E, Thiene G. Classification and histological, immunohistochemical, and molecular diagnosis of inflammatory myocardial disease. *Heart failure reviews.* 2013;18(6):673-681.
17. Cebelin M, Hirsch C. Human stress cardiomyopathy: Myocardial lesions in victims of homicidal assaults without internal injuries. *Hum Pathol.* 1980;11 (2):123-132.
18. Berman M, Ali A, Ashley E, et al. Is stress cardiomyopathy the underlying cause of ventricular dysfunction associated with brain death? *J Heart Lung Transplant.* 2010;29(9):957-965.

19. Wang AR, Dean SA, Grebe SK, Hood IC. Fatal Catecholamine-Induced Cardiotoxicity Associated with Pheochromocytoma: Report of a Postpartum Case and Review of the Literature. *Acad Forensic Pathol.* 2016;6(2):315-324.
20. Frustaci A, Loperfido F, Gentiloni N, Caldarulo M, Morgante E, Russo MA. Catecholamine-induced cardiomyopathy in multiple endocrine neoplasia. A histologic, ultrastructural, and biochemical study. *Chest.* 1991;99(2):382-385.
21. Van Vliet PD, Burchell HB, Titus JL. Focal Myocarditis Associated with Pheochromocytoma. *N Engl J Med.* 1966;274(20):1102-1108.
22. Leone O, Angelini A, Bruneval P, Potena L. *The Pathology of Cardiac Transplantation: A clinical and pathological perspective.* Springer; 2016.
23. Pine DS, Tierney L, Jr. Clinical problem-solving. A stressful interaction. *N Engl J Med.* 1996;334(23):1530-1534.
24. Lindsay J, Paixao A, Chao T, Pichard AD. Pathogenesis of the Takotsubo syndrome: a unifying hypothesis. *Am J Cardiol.* 2010;106(9):1360-1363.
25. Indorato F, Bartoloni G. Post-mortem Takotsubo cardiomyopathy diagnosis: the challenge is open! *Forensic Sci Med Pathol.* 2016;12(2):227-228.
26. Angelini P. Do pathologists agree on how to diagnose takotsubo cardiomyopathy? *Forensic Sci Med Pathol.* 2016;12(2):226.
27. Indorato F, Akashi YJ, Rossitto C, Raffino C, Bartoloni G. Takotsubo cardiomyopathy associated with rupture of the left ventricular apex: assessment of histopathological features of a fatal case and literature review. *Forensic Sci Med Pathol.* 2015;11(4):577-583.

28. Tran K, Milne N, Duhig E, Altman M. Inverted Takotsubo cardiomyopathy-- clinicopathologic correlation. *Am J Forensic Med Pathol.* 2013;34(3):217-221.
29. Savall F, Dedouit F, Telmon N, Rouge D. Takotsubo cardiomyopathy in a jeweler following a hold-up: forensic considerations. *J Forensic Sci.* 2013;58(6):1648-1650.
30. Toni C, Iannaccone F, Chella P, Basolo F, Pucci A. Sudden death in a case of recurrent Takotsubo syndrome. *Forensic Sci Med Pathol.* 2019;15(4):595-597.
31. Akashi YJ, Goldstein DS, Barbaro G, Ueyama T. Takotsubo cardiomyopathy: a new form of acute, reversible heart failure. *Circulation.* 2008;118(25):2754-2762.
32. Assuncao LA, Grams AC, Ribeiro CS, Magalhaes T. "Broken heart" syndrome in an elder abuse case: forensic considerations from the case. *Forensic Sci Med Pathol.* 2011;7(2):217-221.
33. Lyon AR, Rees PS, Prasad S, Poole-Wilson PA, Harding SE. Stress (Takotsubo) cardiomyopathy--a novel pathophysiological hypothesis to explain catecholamine-induced acute myocardial stunning. *Nat Clin Pract Cardiovasc Med.* 2008;5(1):22-29.
34. Boland TA, Lee VH, Bleck TP. Stress-induced cardiomyopathy. *Crit Care Med.* 2015;43(3):686-693.
35. Akashi YJ, Nef HM, Mollmann H, Ueyama T. Stress cardiomyopathy. *Annu Rev Med.* 2010;61:271-286.
36. Sacha J, Maselko J, Wester A, Szudrowicz Z, Pluta W. Left Ventricular Apical Rupture Caused by Takotsubo Cardiomyopathy--Comprehensive Pathological Heart Investigation. *Circ J.* 2007;71:982-985.
37. Wittstein IS, Thiemann DR, Lima JA, et al. Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med.* 2005;352(6):539-548.

38. Gubbi S, Nazari MA, Taieb D, Klubo-Gwiezdzinska J, Pacak K. Catecholamine physiology and its implications in patients with COVID-19. *Lancet Diabetes Endocrinol.* 2020;8(12):978-986.
39. Al Ali AM, Straatman LP, Allard MF, Ignaszewski AP. Eosinophilic myocarditis: case series and review of literature. *Can J Cardiol.* 2006;22(14):1233-1237.
40. Burke AP, Saenger J, Mullick F, Virmani R. Hypersensitivity myocarditis. *Arch Pathol Lab Med.* 1991;115(8):764-769.
41. Verma AK, Lavine KJ, Lin CY. Myocarditis after Covid-19 mRNA Vaccination. *N Engl J Med.* 2021;385(14):1332-1334.
42. Aronoff SC, Hall A, Del Vecchio MT. The Natural History of Severe Acute Respiratory Syndrome Coronavirus 2-Related Multisystem Inflammatory Syndrome in Children: A Systematic Review. *J Pediatric Infect Dis Soc.* 2020;9(6):746-751.

FIGURE LEGENDS

Figure 1: Case A, Heart: Confluent areas of ischemia. Hematoxylin and Eosin stain (H&E), 100X.

Figure 2: Case A, Heart. Coagulative and contraction band necrosis. H&E 200X.

Figure 3: Case A, Heart: Subepicardial fibrosis. This appears older than the timing of the first vaccine dose. This is a possible arrhythmogenic cardiomyopathy, but its appearance is more consistent with healed ischemia or inflammation. H&E 40X.

Figure 4: Case A, Heart. Confluent areas of ischemia with contraction bands and coagulative myocytolysis. 200X.

Figure 5: Case B, Heart: Hypereosinophilic myocytes, contraction band necrosis and coagulative myocytolysis H&E 100X. Inset: The infiltrate is predominantly neutrophilic. H&E 400X.

Figure 6: Case B, Heart. Subepicardial coagulative myocytolysis/contraction band necrosis. H&E 100X.


Figures 7: Case B, Heart. Perivascular inflammation. H&E 200X.


Table 1. Summary of Clinical and Autopsy findings.


Patient	Heart Gross	Microscopic and Molecular
<p>Teenage Boy A, BMI=21. History of attention deficit/hyperactivity syndrome.</p>	<p>280 gms, normal</p>	<p>There was global myocardial injury with areas of coagulative myocytolysis and contraction bands, with a perivascular pattern of inflammation consisting of predominantly neutrophils with histiocytes, scant lymphocytes, and occasional eosinophils (Figures 1-4, Supplemental Figures 1-2). In some sections, the myocardial injury was predominantly subepicardial, while in other sections it was patchy and transmural. In the posterior wall, there was subepicardial/transmural fibrous scar, without fatty replacement. There were no acute or organizing thrombi. The overall pattern of injury was consistent with “stress cardiomyopathy” with contraction bands and a neutrophilic/histiocytic infiltrate.</p> <p>PCR tissue testing performed by the CDC on heart and lung found no molecular evidence of SARS-CoV-2 infection.</p> <p>Molecular testing on postmortem blood detected two variants of uncertain significance: DOLK (c.1257C>G (p.Ile419Met) heterozygous) and MAP2K2 (c.581-3C>T (Intronic) heterozygous).</p>
<p>Teenage Boy B, BMI=30 with obesity</p>	<p>520 grams with biventricular dilatation and marked pulmonary edema (combined lung weight=1481 grams).</p>	<p>There was global myocardial injury similar to that seen above, but with more widespread transmural ischemic changes and more interstitial inflammation, again with a predominant neutrophil component with histiocytes and scant lymphocytes (Figure 5-7, Supplemental Figures 3-4). Several sections had transmural, confluent areas of</p>


		hypereosinophilic myocytes, confluent areas of contraction bands apart from any inflammation, and florid neutrophilic inflammation with some histiocytes. In this case, a subepicardial distribution of injury was not seen. There were no acute or organizing thrombi. PCR tissue testing performed by the CDC on heart and lung found no molecular evidence of SARS-CoV-2 infection.
--	--	--


Abbreviations: BMI, body mass index; CDC, Centers for Disease Control and Prevention; PCR, polymerase chain reaction.

















Supplemental Digital Content. The Supplemental Digital Content was not copyedited by Archives of Pathology & Laboratory Medicine.

Figure 1: Case A, Heart: Confluent contraction band necrosis/coagulative myocytolysis, with a predominantly neutrophilic inflammatory infiltrate with histiocytes. Hematoxylin and Eosin stain (H&E), 100X




Figure 2: Case A, Heart. Interstitial inflammation adjacent to fibrosis. H&E 200X





Figure 3: Case B, Heart. Confluent areas of ischemia with coagulative myocytolysis and contraction band necrosis. H&E 200X



Figures 4: Case B, Heart. Perivascular inflammation. H&E 200X

